




Table 1. Properties of Galaxies

Name	E(B–V)	(m–M) _o	M _V	[Fe/H]	ref.	size ⁵
Pegasus	0.03	24.9	–12.4	–1.1	1	5.0' × 2.7'
	0.15	24.40	–12.3	–1.4	2	
DDO 210	0.03	24.89	–10.9	–1.9	3	2.2' × 1.1'
Tucana	0.0	24.69	–9.3	–1.8	4	2.9' × 1.2'

¹Aparicio 1994²Gallagher et al. 1998³Lee et al. 1999⁴Saviane et al. 1996⁵Taken from the NASA/IPAC Extragalactic Database.




Table 2. Adopted coefficients for I and R–I

numb. stars	Zero point	Color term	Extinction
I	α_i	β_i	k_i
126	-1.753 ± 0.090	0.0237 ± 0.0019	0.30 ± 0.06
R–I	α_{ri}	β_{ri}	k_{ri}
140	-0.770 ± 0.041	0.9958 ± 0.0047	0.00 ± 0.03




Table 3. Journal of observations

Dates	filter	time (s)	FWHM('')	airmass
Pegasus				
10 Aug 99	R	900	1.4	1.454
10 Aug 99	I	600	1.1	1.429
10 Aug 99	TiO	2×1200	1.3	1.415
10 Aug 99	CN	2×1200	1.5	1.434
13 Aug 99	R	900	1.0	1.535
13 Aug 99	I	600	1.1	1.488
13 Aug 99	TiO	1200	1.1	1.462
13 Aug 99	CN	1200	1.1	1.429
DDO 210				
9 Aug 99	I	2×300	1.1	1.048
9 Aug 99	R	2×400	1.2	1.054
9 Aug 99	TiO	2×1200	1.3	1.078
9 Aug 99	CN	2×1200	1.4	1.097
13 Aug 99	R	400	1.1	1.471
13 Aug 99	I	300	1.1	1.439
13 Aug 99	TiO	1200	1.3	1.400
13 Aug 99	CN	1200	1.3	1.331
Tucana				
9 Aug 99	R	2×400	1.3	1.212

Table 3—Continued

Dates	filter	time (s)	FWHM('')	airmass
9 Aug 99	I	2×300	1.2	1.215
9 Aug 99	TiO	2×900	1.4	1.228
9 Aug 99	CN	2×900	1.4	1.257
10 Aug 99	R	400	1.4	1.239
10 Aug 99	I	300	1.1	1.232
10 Aug 99	TiO	900	1.1	1.226
10 Aug 99	CN	900	1.2	1.227




Table 4. C stars in Pegasus

id	RA	Dec	I	σ_I	R–I	σ_{R-I}	CN–TiO	σ_{CN-TiO}
C01	23:28:50.01	14:42:48.5	19.854	0.035	1.316	0.053	0.369	0.085
C02	23:28:35.84	14:44:19.2	19.954	0.032	1.188	0.047	0.408	0.055
C03	23:28:24.79	14:46:31.0	20.008	0.030	0.988	0.041	0.340	0.063
C04	23:28:40.66	14:43:40.1	20.112	0.032	0.956	0.044	0.352	0.066
C05	23:28:48.04	14:41:21.3	20.048	0.033	1.157	0.047	0.180	0.070
C06	23:28:35.89	14:44:47.1	20.200	0.035	1.069	0.049	0.522	0.077
C07	23:28:25.38	14:45:04.3	20.189	0.032	1.081	0.044	0.486	0.075
C08	23:28:34.03	14:44:38.0	20.300	0.035	1.079	0.050	0.301	0.070
C09	23:28:27.65	14:45:14.2	20.326	0.037	1.235	0.056	0.569	0.101
C10	23:28:30.43	14:44:07.6	20.382	0.037	0.969	0.050	0.423	0.085
C11	23:28:44.24	14:43:11.0	20.431	0.043	1.181	0.064	1.073	0.189
C12	23:28:39.54	14:43:16.4	20.422	0.039	1.069	0.054	0.391	0.083
C13	23:28:32.61	14:44:34.4	20.363	0.037	1.365	0.058	0.346	0.081
C14	23:28:29.01	14:45:56.7	20.453	0.040	0.948	0.054	0.330	0.086
C15	23:28:47.19	14:42:58.5	20.420	0.042	1.061	0.060	0.301	0.095
C16	23:28:31.59	14:44:03.7	20.514	0.040	0.968	0.055	0.335	0.086
C17	23:28:41.45	14:43:46.3	20.427	0.039	1.518	0.068	0.565	0.094
C18	23:28:37.35	14:43:49.8	20.524	0.043	1.139	0.063	0.392	0.081
C19	23:28:43.80	14:44:07.0	20.484	0.042	1.087	0.064	0.266	0.095
C20	23:28:32.92	14:44:19.8	20.542	0.039	1.014	0.056	0.398	0.092

Table 4—Continued

id	RA	Dec	I	σ_I	R–I	σ_{R-I}	CN–TiO	σ_{CN-TiO}
C21	23:28:34.35	14:45:23.0	20.481	0.042	1.140	0.060	0.343	0.086
C22	23:28:32.29	14:46:21.5	20.458	0.040	1.376	0.066	0.363	0.093
C23	23:28:41.11	14:44:10.8	20.553	0.043	1.041	0.058	0.731	0.129
C24	23:28:40.07	14:43:41.4	20.558	0.044	1.180	0.067	0.281	0.098
C25	23:28:36.45	14:44:10.6	20.503	0.041	1.443	0.070	0.330	0.092
C26	23:28:38.24	14:43:26.9	20.630	0.048	0.974	0.067	0.279	0.099
C27	23:28:36.58	14:45:08.0	20.664	0.051	1.070	0.071	0.420	0.095
C28	23:28:18.17	14:45:47.7	20.617	0.055	1.101	0.078	0.250	0.124
C29	23:28:23.18	14:45:36.0	20.781	0.058	1.046	0.078	0.450	0.118
C30	23:28:40.61	14:46:13.6	20.790	0.057	0.981	0.076	0.313	0.110
C31	23:28:36.37	14:44:10.0	20.699	0.048	1.497	0.086	0.247	0.102
C32	23:28:36.92	14:42:58.8	20.736	0.051	1.115	0.072	0.310	0.105
C33	23:28:42.56	14:45:03.2	20.853	0.057	0.999	0.080	0.434	0.125
C34	23:28:40.94	14:42:49.7	20.805	0.054	1.247	0.082	0.505	0.121
C35	23:28:42.73	14:44:04.4	20.810	0.057	1.309	0.092	0.644	0.142
C36	23:28:35.58	14:42:16.7	20.918	0.060	0.959	0.080	0.248	0.120
C37	23:28:36.63	14:44:55.2	20.948	0.066	1.204	0.097	0.257	0.110
C38	23:28:47.35	14:41:21.9	20.874	0.071	1.054	0.096	0.213	0.150
C39	23:28:33.74	14:46:37.2	20.886	0.058	1.040	0.079	0.296	0.143
C40	23:28:37.10	14:45:06.4	20.991	0.064	1.049	0.091	0.676	0.153

Table 4—Continued

id	RA	Dec	I	σ_I	R-I	σ_{R-I}	CN-TiO	σ_{CN-TiO}
----	----	-----	---	------------	-----	----------------	--------	-------------------





Table 5. C stars in DDO 210

id	RA	Dec	I	σ_I	R–I	σ_{R-I}	CN–TiO	σ_{CN-TiO}
C01	20:46:50.47	–12:50:24.0	20.020	0.030	1.101	0.051	0.489	0.066
C02	20:46:39.22	–12:50:31.8	20.142	0.039	0.992	0.063	0.313	0.085
C03	20:46:52.92	–12:50:26.2	20.212	0.033	0.933	0.053	0.218	0.070






Table 6. C/M ratios and photometric characteristics

Galaxy	E(R-I)	N_C	N_M	$N_{M_{fg}}$	C/M	$\langle I \rangle$	$\langle R - I \rangle$
Pegasus	0.02	40	135	55	0.73	20.51	1.08
Pegasus	0.12	30	84	44	0.68	20.48	1.14
DDO 210	0.02	3	158	158		20.12	1.01
Tucana	0.00	0	85	85			



Carbon Star Survey of Local Group Dwarf Galaxies. II. Pegasus, DDO 210 and Tucana

Paolo Battinelli¹

Osservatorio Astronomico di Roma, viale del Parco Mellini 84, I-00136 Roma Italy

battinel@oarhp1.rm.astro.it

and

Serge Demers¹

Département de Physique Université de Montréal, Montreal, Qc H3C 3J7, Canada

demers@astro.umontreal.ca

Received _____; accepted _____

accepted by Astronomical Journal (October 2000 Issue)

¹Visiting Astronomer, Cerro Tololo Inter-American Observatory. CTIO is operated by AURA, Inc. under contract to the National Science Foundation.

ABSTRACT

We present the latest results of our ongoing four filter photometric survey of C stars in Local Group dwarf irregular galaxies. Observations of the two low luminosity dwarf irregular galaxies, Pegasus and DDO 210, revealed respectively 40 and 3 C stars, assuming that the reddening of Pegasus is negligible. No C stars were identified in Tucana. Our observations permit the estimation of the CMD contamination by foreground M dwarf thus yielding reliable C/M ratios. Our R, I photometry of the C stars cannot be used to solve the extinction controversy toward Pegasus. The three C stars in DDO 210 are quite bright when compared to C star populations in dwarf galaxies. A larger fainter population in that galaxy seems however improbable. The statistics of C stars, currently on hand for dwarf galaxies, show a well-defined trend with the absolute magnitude of galaxies.

Subject headings: galaxies: individual (Pegasus) (Tucana) (DDO 210) — galaxies: stellar content — stars: carbon

1. INTRODUCTION

Albert et al. (2000, Paper I) have shown the need of an homogeneous set of C stars data for nearby galaxies if one intends to compare carbon star statistics of Local Group galaxies. A flaw of previously published surveys is the very limited area surveyed in each galaxy, thus requiring ill defined extrapolation to assess the global numbers of C or M stars. For example, Cook, Aaronson & Norris (1986) surveyed only two $1.5' \times 2.5'$ areas in NGC 6822 and IC 1613, representing a few percents of the total area of each galaxy. 320×512 pixels CCDs used in the 1980's yielded few C star candidates in external galaxies. For this reason the published statistics are somewhat uncertain. Pritchett et al. (1987) estimated

the C/M ratio of NGC 55 from 12 C stars and 9 M stars for which two were assumed to be foreground dwarfs. Exception to this rule do fortunately exist. Dwarf spheroidal galaxies such as Fornax, Leo I or Phoenix have been entirely surveyed.

Furthermore, different authors have adopted different color limit for the selection of M stars, see Table 1 of Groenewegen (1999). Thus the published C/M ratios are based on statistics for stars later than M0, M3 or M5 making the comparison between galaxies rather difficult. Having the homogeneous survey as a goal, we are then pursuing a systematic survey of Local Group dwarf galaxies to acquire uniform data. This second paper deals with two dwarf irregular galaxies and one dwarf spheroidal galaxy. Table 1 presents the currently known parameters for each galaxy. The size of galaxies is normally based on their bright population. Faint halo of nearby galaxies have been found to extend much farther than their Holmberg radius, see for example the case of DDO 187 investigated by Aparicio, Tikhonov & Karachentsev (2000) and IC 1613 (Paper I).

Pegasus (DDO 216) is among the faint dwarf irregular galaxies of the Local Group. Even though it was discovered in the 1950’s, it was not recognized as a nearby galaxy and possibly a member of the Local Group until Fisher & Tully (1975) detected its HI and measured its radial velocity. Pegasus’ distance has however been somewhat ill defined since its recognition as a possible Local Group member. The first CCD investigation of Pegasus, done by Hoessel & Mould (1982) sets the galaxy well outside the Local Group, at a distance of 1.7 Mpc. A few years later, Hoessel et al. (1990) confirmed this distance using a few newly identified Cepheids. Pegasus’ distance was cut by a factor two by Aparicio (1994) who based his findings on the apparent I magnitude of the tip of the red giant branch. The distance of Pegasus was further reduced by Gallagher et al. (1998) who concluded that Pegasus is metal poor and suffers an unusually large amount of extinction (for an object at $b = -43^\circ$) estimated to be $A_V = 0.47$. As we can see however in Table 1, the adopted

distance has little effect on the calculated absolute magnitude of Pegasus. We shall see that the number of C stars in Pegasus is a strong function of its adopted color excess because our criterion is based on the (R–I) color of the candidates. Hoessel & Mould (1982) have identified 21 stars in Pegasus with $(B-V) > 2.0$. They concluded that these stars, with a $\langle I \rangle = 21.32$, must be carbon stars.

EDITOR: PLACE TABLE ?? HERE.

DDO 210, sometimes called the Aquarius system, is a faint dwarf irregular galaxy which kept its original name from the catalog of dwarf galaxies published by van den Bergh (1959). Its membership to the Local Group was deduced from its radial velocity obtained from HI observations by Lo, Sargent & Young (1993) but only recently a reliable distance was determined by Lee et al. (1999).

Tucana is one of the newly identified members of the ever increasing Local Group family. Listed in the Southern Galaxy Catalog (Corwin et al. 1985), this galaxy was overlooked until Lavery (1990) re-discovered it. Saviane et al. (1996) estimated its distance to be 870 kpc. Tucana is a dwarf spheroidal galaxy of type dE5, according to Lavery and Mighell (1992).

2. OBSERVATIONS

The observations presented in this paper were obtained at the CTIO 1.5 m telescope with a 2048×2048 Tek CCD during a five night run in August 1999. The telescope was employed at the f/13.5 focus, yielding a pixel size of 0.24'' and a field of view of $8.2' \times 8.2'$. This field is deemed suitable for the small galaxies under investigation. It further allows to evaluate the foreground contamination. To photometrically identify C

stars, we employed the technique used in Paper I and described by Brewer et al. (1995). Standard Kron-Cousins R and I filters are used along with CN and TiO interference filters respectively centered at 810 nm and 770 nm. Both filters have a width of 30 nm.

Sky flats were obtained each night through each filters. Calibration to the standard R, I system was done using Landolt’s (1992) equatorial standards observed during the course of the night. Several equatorial fields were observed along with the T Phe field located at declination of -46° . Extinction coefficients and transformation equations were obtained by multilinear regressions. The technique used is similar to the one detailed by Grondin et al. (1990). Once the color term of the magnitude transformation (β_i) and the slope of the color transformation (β_{ri}) were determined for each night, we combined the standards for the five nights to determine one set of coefficients. This set, presented in Table 2, is used to transform all the data. The number of standard stars used in each solution is given in the first column of Table 2.

EDITOR: PLACE TABLE ?? HERE.

We have also investigated the possible need of an atmospheric extinction coefficient for the (CN–TiO) color index. During the course of our run, we observed continuously, during nearly six hours, one of our program galaxy Sgr DIG. The CN and TiO observations were analyzed in the same way than the data presented in this paper. The slopes of the instrumental magnitudes versus airmasses for CN and TiO are both identical and equal to 0.39 ± 0.05 . We thus conclude that there is no need to take into account the atmospheric extinction for the index (CN–TiO).

After the standard prereduction of trimming, bias subtraction, and sky flat-fielding, the photometric reductions were done by fitting model point-spread functions (PSFs) using DAOPHOT/ALLSTAR/ALLFRAME series of programs (Stetson 1987, 1994) in the

following way: we combine, using MONTAGE2, all the images of the target irrespective of the filter to produce a deep image devoid of cosmic rays. ALLSTAR was then used on this deep image to derive a list of stellar images and produce a second image where the stars, found in the first pass, are removed. This subtracted image is also processed through ALLSTAR to find faint stars missed in the first pass. The second list of stars is added to the first one. The final list is then used for the analysis of the individual frames using ALLFRAME. This program fits model PSFs to stellar objects in all the frames simultaneously.

The exposure times were selected to provide a $S/N = 20$ for stars whose magnitude is one magnitude fainter than the expected median magnitude of the C stars. We set, from previous experience, the length of the CN and TiO exposures at least three times the I exposures. The journal of the observations presented here, is given in Table 3. In the I wave band, C stars are among the brightest stars in dwarf galaxies. Thus, we do not need to explore several magnitudes below the red giant tip to observe them all. For this reason, we can survey Local Group galaxies with a medium size telescope.

EDITOR: PLACE TABLE ?? HERE.

2.1. *Comparison with published photometry*

Aparicio & Gallart (1995) have published VRI photometry of Pegasus. The file of photometry was kindly provided by Dr. Aparicio. We are thus in position to compare our magnitudes and colors (BD) with theirs (AG). We were able to match some 1200 stars common to both studies. We compute the following differences.

$$\Delta_i = I_{BD} - I_{AG} = -0.018 \pm 0.002$$

$$\Delta_{ri} = (R - I)_{BD} - (R - I)_{AG} = -0.043 \pm 0.002$$

The color difference is reduced to -0.007 if one compares only stars with $(R-I) > 0.7$. Our magnitudes are thus in excellent agreement with those of Aparicio & Gallart but our overall colors may show a slight color effect relative to their colors. . The Pegasus photometry of Hoessel & Mould (1982) is not in the same photometric system than ours, the magnitude comparison would be quite laborious. It is also nearly impossible to identify the stars from their published small scale photograph.

A similar test can also be done with the short table published by Lee et al. (1999). They, however, do not quote uncertainties for their magnitudes. Comparison of I magnitude of 26 stars, common to both investigations, yields $\Delta_i = -0.02$. A list of Tucana photometry was also published by Saviane et al. (1996). Similarly to the DDO 210 data, no uncertainties are quoted. Furthermore most of their I magnitudes are quite faint and not suitable to be compared with our data. We find, from the comparison of 57 stars, $\Delta_i = -0.05$. These three comparisons give us confidence in our magnitudes and colors.

3. RESULTS

3.1. *Color excess and extinction in our photometric system*

Since we used Kron-Cousins R and I filters, it is necessary to transform published extinctions and color excesses into that system. The relation $E(R-I) = 0.82E(B-V)$ established by Schmidt (1976) holds for the Johnson's R and I filters. However, from Fernie's (1983) relationship between the Kron-Cousins (KC) (R-I) and the Johnson (R-I) for red stars, one sees that the $E(R-I)$ color excess in both systems are nearly identical. We then take Schmidt's relation to calculate the reddening and adopt the relationship between the extinction $A_{I_{KC}}$ and the color excess from Dean et al. (1978): $A_{I_{KC}} = 2.4E(R-I)$.

The interstellar extinction has very little effect on the (CN – TiO) color index because the two wavelengths are near each other. We calculate that $E(\text{CN} - \text{TiO})$ corresponds to 4% of the $E(\text{B}-\text{V})$, thus quite negligible for the galaxies under study. For extreme cases, such as IC 10 which has nearly one magnitude of reddening, one would have to take into account the (CN – TiO) reddening.

EDITOR: PLACE FIGURE 1 HERE.

3.2. *Selection of C and M stars*

We adopt, for the distinction between C and M stars, the same criteria employed in Paper I. The identification of C and M stars is based on the location of the stars in the color-color diagram. The two zones are drawn following the criteria of Paper I. The $R-I = 0.90$ blue limit follows Brewer et al. (1995) original definition. It applies however to a reddening free color-color diagram. The (CN–TiO) index is not expected to vary continuously from M giants to C stars. Carbon suddenly have bands not seen in M stars. The intermediate S stars, with weak CN and TiO bands, are very rare. They will have a (CN–TiO) even more negative than the M stars. Furthermore, C stars are not expected to define a narrow region in the color-color diagram (Cook & Aaronson 1989). This explains why a larger scatter is observed for C stars than for M stars.

The color-color diagram of Pegasus is shown in Figure 1. Here we plot only stars for which $\sigma_{R-I} < 0.1$. This essentially corresponds to stars brighter than $I \approx 21.25$ and with $\sigma_{\text{CN-TiO}} < 0.16$. Relaxing this condition would increase the number of fainter C star candidates but we believe that the reality of their nature would be questionable.

Our observations reveal that there are some forty C stars in Pegasus, three C stars in DDO 210 and none in Tucana. The color-color diagrams obtained for DDO 210 and

Tucana are presented respectively in Figure 2 and Figure 3. The controversy relative to the extinction toward Pegasus compels us to take into account the possibility that its extinction could be unusually high. We discuss this alternative in the next section.

EDITOR: PLACE FIGURE 2 HERE.

EDITOR: PLACE FIGURE 3 HERE.

EDITOR: PLACE TABLE ?? HERE.

The 40 C star candidates in Pegasus, with $(R-I) > 0.92$ are listed in Table 4. The J2000 equatorial coordinates were established using ESO’s Skycat facility and the IRAF tasks CCMAP and CCTRAN. I magnitudes and colors, along with their uncertainties, are presented. Note that this set of C stars is selected by assuming that $E(R-I) = 0.02$. If the color excess is larger, then one would have to move the blue limit of $(R-I)$ to a redder value. Similar data are presented for the three C stars found in DDO 210 in Table 5.

EDITOR: PLACE TABLE ?? HERE.

4. DISCUSSION

4.1. *The evaluation of foreground M stars*

The galaxies selected for our CTIO observations are intentionally small for the CCD field. We can then use the distribution of M stars in the periphery to evaluate the foreground contribution to the total number of M stars. The worst case for this evaluation

is Pegasus because its C star distribution occupies about 70% of the CCD field. C and M stars of the Pegasus field are plotted on Figure 4. The galaxy extends diagonally across the field. We select the two corner areas, defined by the dotted lines, as representative of the foreground M star population. The number of M stars, to be compared with the number of C stars, is given in Table 6 where $N_{M_{fg}}$ refers to the number of foreground stars to be deleted from the total number of M stars, thus yielding a true C/M ratio. For DDO 210, which is aligned E – W, the foreground M stars are evaluated from strips north and south of the galaxy.

EDITOR: PLACE FIGURE 4 HERE.

EDITOR: PLACE FIGURE 5 HERE.

EDITOR: PLACE TABLE ?? HERE.

The magnitude distribution of M stars in the Pegasus field is compared to the distribution of the foreground M stars in the upper panel of Figure 5. Essentially, all M stars brighter than $I \approx 19.6$ are foreground stars. The lower panel of Figure 5 reveals that there is no M stars in Pegasus redder than $R-I = 1.6$, corresponding to spectral type M5 (Thé et al. 1984). The same limit was found in IC 1613 (Paper I).

4.2. *The number of C stars and the M_V of the parent galaxy*

The number of C stars in a galaxy appears to be function of its mass and also of its metal abundance. Such relationships have been presented recently by Groenewegen (1999). We present, in the top panel of Figure 6, what we believe to be the most reliable $\log(N_c)$ vs

M_V relationship. Only galaxies which have been fully surveyed are presented, this condition excludes the Magellanic Clouds. Dwarf spheroidals with blue C stars are excluded: only Fornax (104 C stars) and Leo I (16 C stars) remain. According to this relation, galaxies brighter than $M_V = -15$ are expected to contain several hundred C stars. Excluding the Magellanic Clouds, at least half dozen Local Group dwarfs are as bright as this magnitude. It will eventually be interesting to compare the number of C stars in galaxies of the same absolute magnitude but of different metallicity. NGC 147 and NGC 185, two companions of M31, offer this possibility.

4.3. *C stars and the extinction toward Pegasus*

It would be useful if the observed photometric properties of the C stars in Pegasus could shed some light on the exact value of the extinction along its line of sight. We present, in Table 6, a summary of the photometric properties of C stars in the program galaxies and the calculated C/M ratios. Two entries for Pegasus corresponding to the two currently assumed color excesses are presented. If the color excess of Pegasus is $E(R-I) = 0.12$, corresponding to $E(B-V) = 0.15$, we have to select its C and M stars among stars with $(R-I) > 1.02$. This shift reduces both the number of C stars and the number of M stars, thus modifying slightly the C/M ratio.

According to Gallagher et al. (1998) Pegasus does not only suffer a substantial amount of extinction but is also metal poorer than previously believed. The median color of C stars is known to be a function of the $[Fe/H]$ of the parent galaxy. From the data of Table 6, we can then investigate how the $\langle (R-I)_o \rangle$ for the two alternatives fits other galaxies. The relevant data, taken from Paper I and this paper, are displayed in the lower panel of Figure 6. Pegasus is represented by the two open squares. One sees that both the high color excess with low metallicity and the low color excess with larger metallicity fit the relation. If the

reddening of Pegasus is high, its number of C stars would be 30, such number would still fit the relation between numbers and absolute magnitudes displayed in the upper panel of Figure 6.

EDITOR: PLACE FIGURE 6 HERE.

The AGB models of Groenewegen & de Jong (1993) show that the mean bolometric magnitudes of C stars in the Galaxy and in the LMC are essentially identical, thus possibly independent of the abundance. Current data, on hand, suggest that the median absolute I magnitude of C stars is fairly constant from galaxy to galaxy and is $M_I \approx -4.6$. Our data from Table 6 along with those assembled in Table 1 lead, for Pegasus, to median magnitudes of -4.4 , for the low reddening, and -4.2 for the high reddening. The high reddening yields marginally too faint C stars, if the universality of $\langle M_I \rangle$ is indeed valid.

Clues, as to the precise extinction toward Pegasus, must then come from independent sources and not from C stars. Krienke & Hodge (1998, 2000) have investigated colors of background galaxies seen in the field of Pegasus. They found that background galaxies suffer little reddening when their colors are compared to mean colors of galaxies as compiled by Buta et al. (1994, 1995). This finding suggests that the color excess of Pegasus is normal for its galactic latitude. It is indeed low and Pegasus does have some 40 C stars.

EDITOR: PLACE FIGURE 7 HERE.

The color-magnitude diagrams of Pegasus and the DDO 210 fields are displayed in Figure 7. Stars with $\sigma_{R-I} < 0.3$ are plotted. We see that we have acquired the first two magnitudes of the giant branch. The C stars nicely fit on the extended giant branch. One notices that DDO 210 is at a lower galactic latitude than Pegasus because of the obvious ridge seen at $R-I = 0.4$.

4.4. DDO 210

C02, one of the three C stars of DDO 210, is located 3.2 arcmin from the center of the galaxy. At the distance of DDO 210, this corresponds to nearly 1 kpc from center of DDO 210. It is however along its major axis and well within the region where the old population of DDO 210 is found (Lee et al. 1999). The three stars of DDO 210 are fairly bright when compared to the ones of Pegasus. Could these three stars actually represent the brightest members of an unobserved large population? At least two facts suggest otherwise. First, if we reasonably relax our acceptance criterion, such as allowing stars with larger color errors to be accepted as C stars, we obviously can increase the number of carbon star candidates in DDO 210. The added candidates however are distributed all over the CCD fields and are much fainter than the three listed in Table 5. They do not represent a slightly fainter population in DDO 210. Second, the number of C stars in dwarf galaxies is roughly a function of their mass or absolute magnitude, as we have shown in Fig. 6. DDO 210 is among the faintest dwarf galaxies and it is therefore expected to have few C stars, possibly 4 C stars according to Fig. 6. Thus the C stars in DDO 210 may be unusually bright. Furthermore, DDO 210 must have very few M giants. Our foreground estimate of M stars is identical with the total number of M stars seen in the field. According to the Groenewegen & de Jong (1993) synthetic AGB models, the bulk of M stars is over one magnitude fainter than the C stars, thus a few undetected fainter ones might be present.

5. CONCLUSION

The identification of C stars in a dwarf galaxy points to the presence of an intermediate age population. We have here presented observations of three low mass galaxies. Their numbers of C stars are rather low and preclude further detailed analysis at the present time. Data for Pegasus and DDO 210 indicate that C stars are certainly not restricted to

the core of these galaxies but extend into their halo. C stars in IC 1613 extend further out than its observed HI (Paper I). The HI maps of DDO 210 and Pegasus produced by Lo et al. (1993) show also that the neutral hydrogen is more concentrated than the C stars. C stars could then be more useful than HI to investigate the kinematics of the periphery of disk galaxies such as M31, NGC 55 or WLM. The universality of the median I magnitude of C stars appears, following the DDO 210 observations, to be far from certain. One would definitively need to confirm this with a galaxy with several dozen C stars. Several luminous Local Group dwarf galaxies need to be surveyed for C stars.

This project is supported financially, in part, by the Natural Sciences and Engineering Research Council of Canada (S. D.).

REFERENCES

- Albert, L., Demers, S., & Kunkel, W. E. 2000, AJ, 119, in press (Paper I)
- Aparicio, A. 1994, ApJ, 437, L27
- Aparicio, A., & Gallart, C. 1995, AJ, 110, 2105
- Aparicio, A., Tikhonov, N., & Karachentsev, I. 2000, AJ, 199, 177
- Brewer, J. P., Richer, H. B., & Crabtree, D. R. 1995, AJ, 109, 2480
- Buta, R., Mitra, S., de Vaucouleurs, G., & Corwin, H. G. Jr. 1994, AJ, 107, 118
- Buta, R., Corwin, H. G., Jr., de Vaucouleurs, G., de Vaucouleurs, A., & Longo, G. 1995, AJ, 109, 517
- Cook, K. H., Aaronson, M., & Norris, J. 1986, ApJ, 305, 634
- Cook, K. H., & Aaronson, M. 1989, AJ, 97, 923
- Corwin Jr. H. G., de Vaucouleurs, G., & de Vaucouleurs, A. 1985, Southern Galaxy Catalog (University of Texas Press, Austin, TX)
- Dean, J. F., Warren, P. R., & Cousins, A. W. J. 1978, MNRAS, 183, 569
- Fernie, J. D. 1983, PASP, 95, 782
- Fisher, J. R., & Tully, R. B. 1975, A&A, 44, 151
- Gallagher, J. S., Tolstoy, E., Dohm-Palmer, R. C., Skillman, E. D., Cole, A. A., Hoessel, J. G., Saha, A., & Mateo, M. 1998, AJ, 115, 1869
- Groenewegen, M. A. T. 1999, in IAU Symposium 191 Asymptotic Giant Branch Stars, T. Le Bertre, A. Lèbre and C. Waelkens, eds. ASP, San Francisco, p535

- Groenewegen, M. A. T. & de Jong, T. 1993, *A&A*, 267, 410
- Grondin, L., Demers, S., Kunkel, W. E., & Irwin, M. J. 1990, *AJ*, 100, 663
- Hoessel, J. G., & Mould, J. R. 1982, *ApJ*, 254, 38
- Hoessel, J. G., Abbot, M. J., Saha, A., Mossman, A. E., & Danielson, G. E. 1990, *AJ*, 100, 1151
- Krienke, K., & Hodge, P. W. 1998, *BAAS*, 24,
- Krienke, K., & Hodge, P. W. 2000, Private communication
- Landolt, A. U. 1992, *AJ*, 104, 340
- Lavery, R. J. 1990, *IAU Circular No.* 5139
- Lavery, R. J., & Mighell, K. J. 1992, *AJ*, 103, 81
- Lee, M. G., Aparicio, A., Tikhonov, N., Byun, Y-I., & Kim, E. 1999, *AJ*, 118, 853
- Lo, K. Y., Sargent, W. L. W., & Young, K. 1993, *AJ*, 106, 507
- Pritchett, C. J., Schade, D., Richer, H. B. , Crabtree, D., & Yee, H. K. C. 1987, *ApJ*, 323, 79
- Saviane, I., Held, E. V., & Piotto, G. 1996, *A&A*, 315, 40
- Schmidt, E. G. 1976, *ApJ*, 203, 466
- Stetson, P. B. 1987, *PASP*, 99, 191
- Stetson, P. B. 1994, *PASP*, 106, 250
- Thé, P. S., Steenman, H. C., & Alcaïno, G. 1984, *A&A*, 132, 385
- van den Bergh, S. 1959, *Publ. David Dunlap Obs.* 2, 147

Fig. 1.— The color-color diagram of the stars in the Pegasus field. The blue limit is drawn at $(R-I) = 0.90$ but should be adjusted to take into account the reddening toward Pegasus.

Fig. 2.— The color-color diagram of the DDO 210 field.

Fig. 3.— The color-color diagram of the Tucana field.

Fig. 4.— The spatial distribution of C stars (big dots) and M stars (crosses) in the field of Pegasus. The two dotted lines delineate the “foreground” area. North on top, East to the left.

Fig. 5.— Comparison between the properties of M stars in the field of Pegasus and the properties of M stars in the foreground. The latter are represented by the shaded histograms. The upper panel shows the magnitude distributions and the lower panel presents the color distributions.

Fig. 6.— In the top panel, a plot of the number of C stars, in fully surveyed dwarfs, versus the M_V of the parent galaxy, shows an obvious trend. In the lower panel, we display the relationship between the intrinsic median color of C stars and the $[Fe/H]$ of the parent galaxy. The two open squares represent Pegasus for $E(R-I) = 0.02$ (left) and 0.12 (right) respectively. Both points fit the trend seen among galaxies.

Fig. 7.— Color-magnitude diagram of Pegasus (upper panel) and DDO 210 (lower panel). C stars are represented by big dots. Stars with color uncertainties > 0.3 were excluded, this explains why the limit is $I \approx 22$.